

Mathematics and Computer Science for Modeling

Unit 4: Calculus

Daniel Sabinasz

based on materials by Jan Tekülve and Daniel Sabinasz

Institut für Neuroinformatik, Ruhr-Universität Bochum

September 29, 2023

Overview

1. Motivation


2. Differentiation

- ▶ Graphical Interpretation
- ▶ Formal Description
- ▶ Rules for Differentiation


3. Exercises

Motivation



Estimating Velocity by Differentiation





The Vehicle's Position





The Vehicle's Position





The Vehicle's Position





The Vehicle's Position





The Vehicle's Position





The Vehicle's Position




The Vehicle's Position



The Vehicle's Position



Exercise 1



1. Calculate the change of position between time 3 and 4. Next, calculate the rate of change of the position (= velocity) between time 3 and 4.
2. Do the same for the time between 3 and 5.
3. Now assume that the position is given as $f(t) = t^2$. Plot that function from time 0 to 3. Calculate the velocity between time 0 and 2. Draw a line through the points $(0, f(0))$ and $(2, f(2))$. How does the slope of the line relate to the velocity? Why? Next, do the same for the velocity between time 1 and 2, then between time 1.5 and 2.
4. Think about what it would mean to calculate the velocity *at* time 2.

1. Motivation

2. Differentiation

- ▶ Graphical Interpretation
- ▶ Formal Description
- ▶ Rules for Differentiation



3. Exercises

A function and its derivative

- ▶ The derivative of a function $f(x)$, denoted $f'(x)$, measures the degree to which $f(x)$ changes when x changes


$$f(x) = x$$

$$f'(x) = 1$$



A function and its derivative


- ▶ The derivative of a function $f(x)$, denoted $f'(x)$, measures the degree to which $f(x)$ changes when x changes
- ▶ $f'(x)$ is the slope of the tangent at x




A function and its derivative

$$f(x) = 0.5$$

$$f'(x) = 0$$




A function and its derivative




Derivative as a Tangent

$$f(x) = \sin(x)$$


$$f'(x) = \cos(x)$$



Derivative as a Tangent



Derivative as a Tangent



Formal Definition

Differentiable Function

- ▶ The **derivative of f at position x_0** , short $f'(x)$, is defined as

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

- ▶ This denotes the value of $\frac{f(x) - f(x_0)}{x - x_0}$ as x gets closer and closer to x_0 .

Formal Definition

Differentiable Function

- ▶ The **derivative of f at position x_0** , short $f'(x)$, is defined as

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

- ▶ This denotes the value of $\frac{f(x) - f(x_0)}{x - x_0}$ as x gets closer and closer to x_0 .
- ▶ f is called **differentiable** if and only if this limit exists.

Formal Definition

Differentiable Function

- ▶ The **derivative of f at position x_0** , short $f'(x)$, is defined as

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

- ▶ This denotes the value of $\frac{f(x) - f(x_0)}{x - x_0}$ as x gets closer and closer to x_0 .
- ▶ f is called **differentiable** if and only if this limit exists.
- ▶ Alternate notations:

$$f'(x) = \frac{df}{dx}(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Differentiation as Limit Example

- ▶ **Statement:** The derivative of $f(x) = x^2$ is $f'(x) = 2x$

Differentiation as Limit Example

- ▶ **Statement:** The derivative of $f(x) = x^2$ is $f'(x) = 2x$
- ▶ Applying the formula

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{x \rightarrow x_0} \frac{x^2 - x_0^2}{x - x_0}$$

Differentiation as Limit Example

- ▶ **Statement:** The derivative of $f(x) = x^2$ is $f'(x) = 2x$
- ▶ Applying the formula

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{x \rightarrow x_0} \frac{x^2 - x_0^2}{x - x_0}$$

- ▶ Simplifying

$$\lim_{x \rightarrow x_0} \frac{(x - x_0)(x + x_0)}{x - x_0} = \lim_{x \rightarrow x_0} \frac{\cancel{(x - x_0)}(x + x_0)}{\cancel{x - x_0}} = \lim_{x \rightarrow x_0} (x + x_0)$$

Differentiation as Limit Example

- ▶ **Statement:** The derivative of $f(x) = x^2$ is $f'(x) = 2x$
- ▶ Applying the formula

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{x \rightarrow x_0} \frac{x^2 - x_0^2}{x - x_0}$$

- ▶ Simplifying

$$\lim_{x \rightarrow x_0} \frac{(x - x_0)(x + x_0)}{x - x_0} = \lim_{x \rightarrow x_0} \frac{\cancel{(x - x_0)}(x + x_0)}{\cancel{x - x_0}} = \lim_{x \rightarrow x_0} (x + x_0)$$

- ▶ Applying the limit:

$$\lim_{x \rightarrow x_0} (x + x_0) = 2x$$

Differentiation is a linear operator

Rules

▶ **Constant Factor**

$$\frac{d}{dx}(af) = a \frac{d}{dx}(f)$$

▶ **Sums**

$$\frac{d}{dx}(f + g) = \frac{d}{dx}(f) + \frac{d}{dx}(g)$$

Example:

$$\frac{d}{dx}(4x^2) = 4 \frac{d}{dx}(x^2) = 4(2x) = 8x$$

Differentiation is a linear operator

Rules

▶ **Constant Factor**

$$\frac{d}{dx}(af) = a \frac{d}{dx}(f)$$

▶ **Sums**

$$\frac{d}{dx}(f + g) = \frac{d}{dx}(f) + \frac{d}{dx}(g)$$

Example:

$$\frac{d}{dx}(4x^2) = 4 \frac{d}{dx}(x^2) = 4(2x) = 8x$$

$$\frac{d}{dx}(4x^2 + x^2) = 4 \frac{d}{dx}(x^2) + \frac{d}{dx}(x^2) = 4(2x) + 2x = 10x$$

Differentiation for Products and Quotients

Rules

▶ **Multiplication**

$$\frac{d}{dx}(fg) = \frac{d}{dx}(f)g + f\frac{d}{dx}(g)$$

▶ **Exponentiation**

$$\frac{d}{dx}(f^n) = n\frac{d}{dx}(f)^{n-1}$$

▶ **Division**

$$\frac{d}{dx}\left(\frac{f}{g}\right) = \frac{\frac{d}{dx}(f)g - f\frac{d}{dx}(g)}{g^2}$$

Examples

► Multiplication

$$\frac{d}{dx}(x^2 \sin(x)) = \frac{d}{dx}(x^2) \sin(x) + x^2 \frac{d}{dx}(\sin(x)) = 2x \sin(x) + x^2 \cos(x)$$

Examples

► Multiplication

$$\frac{d}{dx}(x^2 \sin(x)) = \frac{d}{dx}(x^2) \sin(x) + x^2 \frac{d}{dx}(\sin(x)) = 2x \sin(x) + x^2 \cos(x)$$

► Division

$$\frac{d}{dx} \left(\frac{1}{x} \right) = \frac{\frac{d}{dx}(1)x - 1 \frac{d}{dx}(x)}{x^2} = \frac{0 - 1}{x^2} = \frac{-1}{x^2}$$

Exponentiation Rule derives from Multiplication Rule

► Example $f'(x^3)$

$$\frac{d}{dx}(x^3) = \frac{d}{dx}(x^2x) = \frac{d}{dx}(x^2)x + x^2 \frac{d}{dx}(x)$$

Exponentiation Rule derives from Multiplication Rule

► Example $f'(x^3)$

$$\begin{aligned}\frac{d}{dx}(x^3) &= \frac{d}{dx}(x^2x) = \frac{d}{dx}(x^2)x + x^2 \frac{d}{dx}(x) \\ &= 2xx + x^2 = 3x^2\end{aligned}$$

Exponentiation Rule derives from Multiplication Rule

- ▶ Example $f'(x^3)$

$$\begin{aligned}\frac{d}{dx}(x^3) &= \frac{d}{dx}(x^2x) = \frac{d}{dx}(x^2)x + x^2 \frac{d}{dx}(x) \\ &= 2xx + x^2 = 3x^2\end{aligned}$$

- ▶ Example $f'(x^4)$

$$\frac{d}{dx}(x^4) = \frac{d}{dx}(x^2x^2) = \frac{d}{dx}(x^2)x^2 + x^2 \frac{d}{dx}(x^2)$$

Exponentiation Rule derives from Multiplication Rule

- Example $f'(x^3)$

$$\begin{aligned}\frac{d}{dx}(x^3) &= \frac{d}{dx}(x^2x) = \frac{d}{dx}(x^2)x + x^2 \frac{d}{dx}(x) \\ &= 2xx + x^2 = 3x^2\end{aligned}$$

- Example $f'(x^4)$

$$\begin{aligned}\frac{d}{dx}(x^4) &= \frac{d}{dx}(x^2x^2) = \frac{d}{dx}(x^2)x^2 + x^2 \frac{d}{dx}(x^2) \\ &= 2xx^2 + x^2 2x = 2x^3 + 2x^3 = 4x^3\end{aligned}$$

Special cases

- ▶ The derivative of

$$f(x) = e^x \text{ is } f'(x) = e^x$$

- ▶ The derivative of

$$f(x) = \ln(x) \text{ is } f'(x) = \frac{1}{x}$$

- ▶ The derivative of

$$f(x) = \sin(x) \text{ is } f'(x) = \cos(x)$$

Composite functions

Chain Rule

- ▶ Function h is a composition of functions g and f

$$h(x) = (g \circ f)(x) = g(f(x))$$

- ▶ If f and g are differentiable, h is also differentiable

$$\frac{d}{dx}(h(x)) = \frac{d}{dx}(g(y)) \frac{d}{dx}(f(x)), \text{ with } y = f(x)$$

- ▶ Verbal rule: **Inner derivative times outer derivative**

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

$$g'(x) = 20x^3 \wedge f'(x) = 7$$

Chain Rule Examples

► $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

$$g'(x) = 20x^3 \wedge f'(x) = 7$$

$$h'(x) = 20(7x + 2)^3 \cdot 7 = 140(7x + 2)^3$$

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

$$g'(x) = 20x^3 \wedge f'(x) = 7$$

$$h'(x) = 20(7x + 2)^3 \cdot 7 = 140(7x + 2)^3$$

▶ $h(x) = e^{5x} = g(f(x))$

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

$$g'(x) = 20x^3 \wedge f'(x) = 7$$

$$h'(x) = 20(7x + 2)^3 \cdot 7 = 140(7x + 2)^3$$

▶ $h(x) = e^{5x} = g(f(x))$

$$g(x) = e^x \wedge f(x) = 5x$$

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

$$g'(x) = 20x^3 \wedge f'(x) = 7$$

$$h'(x) = 20(7x + 2)^3 \cdot 7 = 140(7x + 2)^3$$

▶ $h(x) = e^{5x} = g(f(x))$

$$g(x) = e^x \wedge f(x) = 5x$$

$$g'(x) = e^x \wedge f'(x) = 5$$

Chain Rule Examples

▶ $h(x) = 5(7x + 2)^4 = g(f(x))$

$$g(x) = 5x^4 \wedge f(x) = 7x + 2$$

$$g'(x) = 20x^3 \wedge f'(x) = 7$$

$$h'(x) = 20(7x + 2)^3 7 = 140(7x + 2)^3$$


▶ $h(x) = e^{5x} = g(f(x))$

$$g(x) = e^x \wedge f(x) = 5x$$


$$g'(x) = e^x \wedge f'(x) = 5$$

$$h'(x) = e^{5x} 5 = 5e^{5x}$$

Finding Local Extrema



Finding Local Extrema



Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

Calculation of Local Extrema

► $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

Calculation of Local Extrema

$$\blacktriangleright f(x) = 4x^2 + 6x$$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

Calculation of Local Extrema

$$\blacktriangleright f(x) = 4x^2 + 6x$$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

$$\iff x = \frac{-6}{8} = \frac{-3}{4}$$

Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

$$\iff x = \frac{-6}{8} = \frac{-3}{4}$$

▶ $f(x) = \sin(x)$

Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

$$\iff x = \frac{-6}{8} = \frac{-3}{4}$$

▶ $f(x) = \sin(x)$

$$f'(x) = \cos(x)$$

Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

$$\iff x = \frac{-6}{8} = \frac{-3}{4}$$

▶ $f(x) = \sin(x)$

$$f'(x) = \cos(x)$$

$$f'(x) = \cos(x) \stackrel{!}{=} 0$$

Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

$$\iff x = \frac{-6}{8} = \frac{-3}{4}$$

▶ $f(x) = \sin(x)$

$$f'(x) = \cos(x)$$

$$f'(x) = \cos(x) \stackrel{!}{=} 0$$

$$\iff x = \cos^{-1}(0)$$

Calculation of Local Extrema

▶ $f(x) = 4x^2 + 6x$

$$f'(x) = 8x + 6$$

$$f'(x) = 8x + 6 \stackrel{!}{=} 0$$

$$\iff 8x = -6$$

$$\iff x = \frac{-6}{8} = \frac{-3}{4}$$

▶ $f(x) = \sin(x)$

$$f'(x) = \cos(x)$$


$$f'(x) = \cos(x) \stackrel{!}{=} 0$$

$$\iff x = \cos^{-1}(0)$$


$$\iff x = 90^\circ = \frac{\pi}{2}, 270^\circ = \frac{3\pi}{2}, \dots$$

Differentiability is not given

$$f(x) = \frac{1}{x}$$



$$f'(x) = \frac{-1}{x^2}$$



Exercise 2

1. Calculate the derivative of the following functions (on a piece of paper)

1.1 $f(x) = 7x^4$

1.2 $g(x) = 2x^4 + 3x^3 + x^2 + 10x + 5$

1.3 $h(x) = 4e^{3x}$

1.4 $i(x) = (12x^2 + 5)3x^3$

1.5 $j(x) = \frac{3x}{\cos(x)}$

- ▶ First think about the rule you need to use
- ▶ Identify the parts of the rule in the equation
- ▶ If possible differentiate individual parts first
- ▶ Apply the rule

2. Find the extreme value of the function $k(x) = 6x^2 + 3x + 2$